arXiv:1110.2999v4 [hep-th] 7 May 2012

A Higher Order GUP with Minimal Length Uncertainty and Maximal Momentum

Pouria Pedram*

Department of Physics, Science and Research Branch,

Islamic Azad University, Tehran, Iran

(Dated: May 8, 2012)

Abstract

We present a higher order Generalized (Gravitational) Uncertainty Principle (GUP) in the form $[X, P] = i\hbar/(1 - \beta P^2)$. This form of GUP is consistent with various proposals of quantum gravity such as string theory, loop quantum gravity, doubly special relativity, and predicts both a minimal length uncertainty and a maximal observable momentum. Here we present two equivalent and exact representations where one is formally self-adjoint and naturally perturbative. We show that the presence of the maximal momentum results in an upper bound on the energy spectrum of the free particle, particle in box, and harmonic oscillator. We then generalize this GUP to D dimensions that will be shown it is noncommutative and find the invariant density of states. This form of GUP modifies the blackbody radiation spectrum at high frequencies and predicts a finite cosmological constant. Although it does not solve the cosmological constant problem, it gives a better estimation with respect to the presence of just the minimal length.

PACS numbers: 04.60.Bc

Keywords: Quantum gravity; Generalized uncertainty principle; Minimal length; Maximal momentum.

---

* p.pedram@srbiau.ac.ir

I. INTRODUCTION

In recent years, there is a great interest to study the effects of the Generalized Uncertainty Principle (GUP) and the Modified Dispersion Relation (MDR) on various quantum mechanical systems. Indeed, the ideas of GUP and MDR arise naturally from various candidates of quantum gravity such as string theory [1–4], loop quantum gravity [5], noncommutative spacetime [6–8], and black holes gedanken experiments [9, 10]. These theories indicate that the Heisenberg uncertainty principle should be modified to incorporate additional constraints in the presence of the gravitational field.

The existence of a minimal length scale of the order of the Planck length $\ell_{Pl} = \sqrt{\frac{G\hbar}{c^3}} \approx 10^{-35}m$ is one of the main outcomes of various GUP proposals where G is Newton's gravitational constant. In fact, beyond the Planck energy scale, the effects of gravity are so important which would result in discreteness of the very spacetime. Notably, the quantum field theory in curved background can be renormalizable by introducing a minimal observable length as an effective cutoff in the ultraviolet domain. Also, in the string-theoretic argument, we can say that the string cannot probe distances smaller than its own length.

The introduction of this idea has drawn much attention in the literature to study the effects of GUP on small scale and large scale systems [11–26]. It is also possible to incorporate the idea of a maximal observable momentum into this scenario. In fact, in doubly special relativity (DSR) theories, we consider the Planck energy (Planck Momentum) as an additional invariant other than the velocity of light [27–29]. Recently, the construction of a perturbative GUP which is consistent with DSR theories is also discussed in Refs. [30–34]. It is also shown that a minimum uncertainty in momentum can arise from curvature, as part of a study that indicated that curvature and noncommutativity can be seen as dual to each other [35].

In this paper, we investigate the effects of a new generalized uncertainty principle to all orders in the Planck length on some quantum mechanical systems. This form of GUP implies the existence of a minimal length uncertainty and a maximal momentum in agreement with various theories of quantum gravity. We present two equivalent exact representations of this modified algebra where the latter is self-adjoint and does not change the nature of the position operator. Moreover, the existence of the maximal momentum is manifest from this representation.

Here, we study the problems of the free particle, particle in a box, and the harmonic oscillator in this scenario and obtain their quantum mechanical and semiclassical energy spectrum which are bounded from above. We also address the maximal localization states, generalization to D dimensions, validity of semiclassical approximation, invariant density of states, cosmological constant, and blackbody radiation in this GUP framework.

II. THE GENERALIZED UNCERTAINTY PRINCIPLE

First let us consider a generalized uncertainty principle proposed by Kempf, Mangano and Mann (KMM) and results in a minimum observable length

$$\Delta X \Delta P \geq \frac{\hbar}{2}\left(1+\beta(\Delta P)^2+\zeta\right), \tag{1}$$

where β is the GUP parameter and ζ is a positive constant that depends on the expectation values of the momentum operator, i.e., $\zeta = \beta\langle P\rangle^2$. We also have $\beta = \beta_0/(M_{Pl}c)^2$ where M_{Pl} is the Planck mass and β_0 is of the order of the unity. It is straightforward to check that the inequality relation (1) implies the existence of a minimum observable length as $(\Delta X)_{min}^{\mathrm{KMM}} = \hbar\sqrt{\beta}$. In one-dimension, the above uncertainty relation can be obtained from the following deformed commutation relation:

$$[X,P] = i\hbar(1+\beta P^2). \tag{2}$$

As KMM have indicated in their seminal paper, we can write X and P in momentum space representation as [7]

$$P\phi(p) = p\,\phi(p), \tag{3}$$

$$X\phi(p) = i\hbar\left(1+\beta p^2\right)\partial_p\phi(p), \tag{4}$$

where X and P are symmetric operators on the dense domain S_∞ with respect to the following scalar product:

$$\langle\psi|\phi\rangle = \int_{-\infty}^{+\infty}\frac{\mathrm{d}p}{1+\beta p^2}\psi^*(p)\phi(p), \tag{5}$$

where $\int_{-\infty}^{+\infty}\frac{\mathrm{d}p}{1+\beta p^2}|p\rangle\langle p| = 1$ and $\langle p|p'\rangle = (1+\beta p^2)\,\delta(p-p')$. With this definition, the commutation relation (2) is exactly satisfied.

Based on the field theory on nonanticommutative superspace, Nouicer has suggested the following higher order GUP which agrees with (2) to the leading order and also predicts a minimal length uncertainty

$$[X, P] = i\hbar \exp\left(\beta P^2\right). \tag{6}$$

This algebra can be satisfied from the following representation of the position and momentum operators:

$$P\phi(p) = p\,\phi(p), \tag{7}$$

$$X\phi(p) = i\hbar \exp\left(\beta p^2\right) \partial_p \phi(p). \tag{8}$$

Now the symmetricity condition of the position operator implies the following modified completeness relation and scalar product

$$\langle\psi|\phi\rangle = \int_{-\infty}^{+\infty} \mathrm{d}p \exp\left(-\beta p^2\right) \psi^*(p)\phi(p), \tag{9}$$

$$\langle p|p'\rangle = \exp\left(\beta p^2\right) \delta(p - p'). \tag{10}$$

Also, the absolutely smallest uncertainty in position is given by $(\Delta X)_{min}^{\text{Nouicer}} = \sqrt{\frac{e}{2}}\hbar\sqrt{\beta}$.

To incorporate the idea of the maximal momentum, Ali, Das and Vagenas have proposed the following modified commutation relation [32–34]

$$[X_i, P_j] = i\hbar\left[\delta_{ij} - \alpha\left(P\delta_{ij} + \frac{P_i P_j}{P}\right) + \alpha^2\left(P^2\delta_{ij} + 3P_i P_j\right)\right], \tag{11}$$

where $\alpha = \alpha_0/M_{Pl}c = \alpha_0\ell_{Pl}/\hbar$ is the GUP parameter, $P^2 = \sum_{j=1}^{3} P_j P_j$, M_{Pl} is the Planck mass, and $M_{Pl}c^2 \sim 10^{19}$GeV is the Planck energy. This form of GUP implies both a minimal length uncertainty and a maximal momentum uncertainty, namely [32]

$$\Delta X \geq (\Delta X)_{min} \approx \alpha_0\ell_{Pl} = \hbar\alpha, \tag{12}$$

$$\Delta P \leq (\Delta P)_{max} \approx \frac{M_{Pl}c}{\alpha_0} = 1/\alpha. \tag{13}$$

The commutation relation (11) is approximately satisfies by the the following representation

$$X_i = x_i, \tag{14}$$

$$P_i = p_i\left(1 - \alpha p + 2\alpha^2 p^2\right), \tag{15}$$

where x_i and p_i obey the usual commutation relations $[x_i, p_j] = i\hbar\delta_{ij}$ and p is the magnitude of $\vec{p}$. Now Eq. (12) implies $\alpha \approx \sqrt{\beta}$. However, this proposal has the following difficulties:

- It is perturbative, i.e., it is only valid for small values of the GUP parameter.

- Although the minimal length uncertainty can be considered in some sense as the minimal length, the maximal momentum uncertainty cannot be interpreted as the maximal momentum in the same way which is required in DSR theories. Indeed Eq. (13) puts an upper bound on the uncertainty of the momentum measurement, not on the value of the observerd momentum.

- It does not imply noncommutative geometry, because $[X_i, X_j] = 0$ [see Eq. (14)].

To overcome these problems, consider the following higher order generalized uncertainty principle (GUP*) which implies both the minimal length uncertainty and the maximal observable momentum

$$[X, P] = \frac{i\hbar}{1 - \beta P^2}. \tag{16}$$

This commutation relation agrees with KMM's and Noucier's proposals to the leading order and contains a singularity at $P^2 = 1/\beta$. This fact shows that the momentum of the particle cannot exceed $1/\sqrt{\beta} \approx 1/\alpha$ which agrees formally with Eq. (13). As stated before, Eqs. (13) and (16) imply two basically different quantities. However, the presence of an upper bound on the momentum properly agrees with DSR theories. As we shall see, the physical observables such as energy and momentum are not only nonsingular, but also are bounded from above.

To satisfy the above commutation relation, we can write the position and momentum operators in the momentum space representation as

$$P\phi(p) = p\,\phi(p), \tag{17}$$

$$X\phi(p) = \frac{i\hbar}{1 - \beta p^2}\partial_p\phi(p). \tag{18}$$

Using the symmetricity condition of the position operator the modified completeness relation and scalar product can be written as

$$\langle\psi|\phi\rangle = \int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \mathrm{d}p \left(1 - \beta p^2\right) \psi^*(p)\phi(p), \tag{19}$$

$$\langle p|p'\rangle = \frac{\delta(p - p')}{1 - \beta p^2}. \tag{20}$$

TABLE I. The minimal length uncertainties and maximal momentums in three GUP frameworks.

	KMM	Nouicer	GUP*
$(\Delta X)_{min}$	$\hbar\sqrt{\beta}$	$\sqrt{\frac{e}{2}}\hbar\sqrt{\beta}$	$\frac{3\sqrt{3}}{4}\hbar\sqrt{\beta}$
P_{max}	–	–	$\frac{1}{\sqrt{\beta}}$

The uncertainty relation that arises from GUP* is given by

$$
\begin{aligned}
(\Delta X)(\Delta P) &\geq \left\langle \frac{\hbar/2}{1-\beta P^2} \right\rangle, \\
&\geq \frac{\hbar}{2}\left(1+\beta\left\langle P^2\right\rangle+\beta^2\left\langle P^4\right\rangle+\beta^3\left\langle P^6\right\rangle+\cdots\right), \\
&\geq \frac{\hbar}{2}\left(1+\beta\left\langle P^2\right\rangle+\beta^2\left\langle P^2\right\rangle^2+\beta^3\left\langle P^2\right\rangle^3+\cdots\right), \\
&\geq \frac{\hbar}{2}\Big(1+\beta\left[(\Delta P)^2+\langle P\rangle^2\right]+\beta^2\left[(\Delta P)^2+\langle P\rangle^2\right]^2+\beta^3\left[(\Delta P)^2+\langle P\rangle^2\right]^3+\cdots\Big), \\
&\geq \frac{\hbar/2}{1-\beta\left[(\Delta P)^2+\langle P\rangle^2\right]},
\end{aligned}
\tag{21}
$$

where we have used the property $\langle P^{2n}\rangle \geq \langle P^2\rangle^n$. In order to find the minimal length uncertainty of this deformed algebra, we consider the physical states for which we have $\langle P\rangle = 0$ and solve the following saturate GUP* for ΔP

$$(\Delta X)(\Delta P) = \frac{\hbar/2}{1-\beta(\Delta P)^2}, \tag{22}$$

which has a minimum at $\Delta P = 1/\sqrt{3\beta}$. So the absolutely smallest uncertainty in position is given by

$$(\Delta X)^*_{min} = \frac{3\sqrt{3}}{4}\hbar\sqrt{\beta}. \tag{23}$$

In Table I, we have compared minimal length uncertainties from various GUP scenarios. These results show that $(\Delta X)^{\text{KMM}}_{min} < (\Delta X)^{\text{Noucier}}_{min} < (\Delta X)^*_{min}$.

III. FUNCTIONAL ANALYSIS OF THE POSITION OPERATOR

The eigenvalue problem for the position operator in the GUP* framework and in the momentum space is given by

$$\frac{i\hbar}{1-\beta p^2}\partial_p\psi_\lambda(p) = \lambda\psi_\lambda(p). \tag{24}$$



FIG. 1. Plotting $\langle\psi_\lambda|\psi_{\lambda'}\rangle$ over $\lambda-\lambda'$ in units of $\hbar\sqrt{\beta}$ for the KMM GUP (blue line) and GUP* (red line).

This equation can be solved to obtain the position eigenvectors

$$\psi_\lambda(p) = c\exp\left(\frac{-i\lambda p}{\hbar}\left(1-\frac{\beta}{3}p^2\right)\right). \tag{25}$$

The eigenfunctions are normalizable

$$1 = cc^*\int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \mathrm{d}p\,\left(1-\beta p^2\right) = \frac{4cc^*}{3\sqrt{\beta}}. \tag{26}$$

Therefore

$$\psi_\lambda(p) = \frac{\sqrt{3\sqrt{\beta}}}{2}\exp\left(\frac{-i\lambda p}{\hbar}\left(1-\frac{\beta}{3}p^2\right)\right). \tag{27}$$

Now we calculate the scalar product of the position eigenstates

$$\begin{aligned}\langle\psi_\lambda|\psi_{\lambda'}\rangle &= \frac{3\sqrt{\beta}}{4}\int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}}\left(1-\beta p^2\right)\exp\left(\frac{i(\lambda-\lambda')p}{\hbar}\left(1-\frac{\beta}{3}p^2\right)\right)\mathrm{d}p,\\ &= \frac{3\hbar\sqrt{\beta}}{2(\lambda-\lambda')}\sin\left(\frac{2(\lambda-\lambda')}{3\hbar\sqrt{\beta}}\right).\end{aligned} \tag{28}$$

Thus, similar to the KMM scenario, the position eigenstates are generally no longer orthogonal. In Fig. 1, we have depicted $\langle\psi_\lambda|\psi_{\lambda'}\rangle$ for the KMM proposal and GUP*. Although this quantity in both models has a same functional form, it is more oscillatory in the KMM framework.

A. Maximal localization states

The maximal localization states $|\psi_\xi^{\mathrm{ML}}\rangle$ are defined with the properties

$$\langle\psi_\xi^{\mathrm{ML}}|X|\psi_\xi^{\mathrm{ML}}\rangle = \xi, \tag{29}$$

and

$$\Delta X_{|\psi_\xi^{\mathrm{ML}}\rangle} = (\Delta X)^*_{min}. \tag{30}$$

These states also satisfy [7]

$$\left(X - \langle X\rangle + \frac{\langle[X,P]\rangle}{2(\Delta P)^2}\left(P - \langle P\rangle\right)\right)|\psi\rangle = 0. \tag{31}$$

To proceed further we need to express $\langle[X,P]\rangle$ in terms of ΔP and $\langle P\rangle$. However, since $\langle[X,P]\rangle$ also depends on $\langle P^4\rangle$, $\langle P^6\rangle$, etc., and these quantities cannot be calculated before specifying $|\psi\rangle$, to first order in the GUP parameter we can use the approximate relation $\langle[X,P]\rangle \simeq i\hbar\left(1+\beta(\Delta P)^2+\beta\langle P\rangle^2\right)$. So, in momentum space, the above equation takes the form

$$\left(\frac{i\hbar}{1-\beta p^2}\frac{\partial}{\partial p} - \langle X\rangle + i\hbar\frac{1+\beta(\Delta P)^2+\beta\langle P\rangle^2}{2(\Delta P)^2}\Big(p-\langle P\rangle\Big)\right)\psi(p) \simeq 0, \tag{32}$$

which has the solution

$$\begin{aligned}\psi(p) \simeq&\mathcal{N}\exp\Bigg[\left(-\frac{i}{\hbar}\langle X\rangle + \frac{1+\beta(\Delta P)^2+\beta\langle P\rangle^2}{2(\Delta P)^2}\langle P\rangle\right)\\ &\times\left(p-\frac{\beta}{3}p^3\right) - \frac{1+\beta(\Delta P)^2+\beta\langle P\rangle^2}{4(\Delta P)^2}\left(p^2-\frac{\beta}{2}p^4\right)\Bigg].\end{aligned} \tag{33}$$

To find the absolutely maximal localization states we need to choose the critical momentum uncertainty $\Delta P = 1/\sqrt{3\beta}$ that gives the minimal length uncertainty and take $\langle P\rangle = 0$, i.e.,

$$\psi_\xi^{\mathrm{ML}}(p) \simeq \mathcal{N}\exp\left[-\frac{i}{\hbar}\xi\left(p-\frac{\beta}{3}p^3\right) - \beta\left(p^2-\frac{\beta}{2}p^4\right)\right], \tag{34}$$

where the normalization factor is given by

$$\begin{aligned}1 &= \mathcal{N}\mathcal{N}^*\int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \mathrm{d}p\left(1-\beta p^2\right)\exp\left(2\beta p^2-\beta^2p^4\right),\\ &= 1.0123\frac{\mathcal{N}^2}{\sqrt{\beta}}.\end{aligned} \tag{35}$$

Note that $\psi_\xi^{\mathrm{ML}}(p)$ exactly satisfies Eq. (29). However, because of the approximation that assumed to find $\psi_\xi^{\mathrm{ML}}(p)$ (34), it approximately obeys relation (30), i.e.,

$$\Delta X_{|\psi_\xi^{\mathrm{ML}}\rangle} = 1.0998(\Delta X)^*_{min}, \tag{36}$$

which shows an error less than 10%. Also, because of the fuzziness of space, these maximal localization states are not mutually orthogonal. It is worth to mention that, in this framework, the expectation value of the kinetic energy operator $P^2/2m$ is finite for both $|\psi_\lambda\rangle$ and $|\psi_\xi^{\mathrm{ML}}\rangle$. Indeed we have

$$\left\langle \psi_\lambda \middle| \frac{P^2}{2m} \middle| \psi_\lambda \right\rangle = \frac{1}{10m\beta}, \tag{37}$$

and

$$\left\langle \psi_\xi^{\mathrm{ML}} \middle| \frac{P^2}{2m} \middle| \psi_\xi^{\mathrm{ML}} \right\rangle = \frac{0.7345}{10m\beta}. \tag{38}$$

These quantities for the KMM proposal are ∞ and $1/2m\beta$, respectively.

To find the quasiposition wave function $\psi(\xi)$, we define

$$\psi(\xi) \equiv \langle \psi_\xi^{\mathrm{ML}} | \psi \rangle, \tag{39}$$

where in the limit $\beta \to 0$ it goes to the ordinary position wave function $\psi(\xi) = \langle \xi | \psi \rangle$. Now the transformation of the wave function in the momentum representation into its counterpart quasiposition wave function is

$$\psi(\xi) = \mathcal{N} \int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \mathrm{d}p \, (1 - \beta p^2) \exp\left[\frac{i}{\hbar} \xi \left(p - \frac{\beta}{3} p^3 \right) - \beta \left(p^2 - \frac{\beta}{2} p^4 \right) \right] \psi(p). \tag{40}$$

This relation shows that similar to the ordinary quantum mechanics and the KMM proposal, the quasiposition wave function of a momentum eigenstate $\psi_{\tilde{p}}(p) = \delta(p - \tilde{p})$ with energy $E = \tilde{p}^2/2m$ is still a plane wave but with a modified dispersion relation

$$\lambda(E) = \frac{2\pi\hbar}{\sqrt{2mE}\left(1 - \frac{2}{3} m\beta E\right)} = \frac{\lambda_{\mathrm{ord}}(E)}{1 - \frac{2}{3} m\beta E}, \tag{41}$$

where $\lambda_{\mathrm{ord}}(E) = 2\pi\hbar/\sqrt{2mE}$ is the wavelength in the absence of GUP. In Fig. 2 we have depicted λ versus mE in various scenarios. Since Eq. (41) is bounded from below, there exists a nonzero minimal wavelength. So the wavelength components smaller than

$$\lambda_0 = 3\pi\hbar\sqrt{\beta} = \frac{3}{4}\pi\lambda_0^{\mathrm{KMM}}, \tag{42}$$

are absent in the Fourier decomposition of the quasiposition wave function of the physical states. Therefore, the maximal energy of a momentum eigenstate is

$$E_{max} = \frac{3}{2m\beta}. \tag{43}$$



FIG. 2. The wavelength of the quasiposition wave function of a momentum eigenstate in ordinary quantum mechanics, KMM's GUP and GUP* for $\beta = 0.2$.

Since the transformation (40) as the generalized Fourier transformation is invertible, the transformation of a quasiposition wave function into a momentum space wave function is given by

$$\psi(p) = \frac{\mathcal{N}^{-1}}{2\pi\hbar} \int_{-\infty}^{+\infty} \mathrm{d}\xi \, \exp\left[\beta\left(p^2 - \frac{\beta}{2}p^4\right)\right] \exp\left[-\frac{i}{\hbar}\xi\left(p - \frac{\beta}{3}p^3\right)\right]\psi(\xi). \tag{44}$$

Now the scalar product of states in terms of the quasiposition wave functions reads

$$\begin{aligned}\langle\phi|\psi\rangle &= \int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \mathrm{d}p \,(1-\beta p^2)\phi^*(p)\psi(p), \\ &= \left(\frac{\mathcal{N}^{-1}}{2\pi\hbar}\right)^2 \int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \mathrm{d}p \, \mathrm{d}\xi \, \mathrm{d}\xi'(1-\beta p^2) \exp\left[2\beta\left(p^2 - \frac{\beta}{2}p^4\right)\right] \\ &\quad \times \exp\left[-\frac{i}{\hbar}(\xi - \xi')\left(p - \frac{\beta}{3}p^3\right)\right]\phi^*(\xi)\psi(\xi'). \end{aligned} \tag{45}$$

IV. HARMONIC OSCILLATOR

In this section, we apply the developed formalism to the case of a linear harmonic oscillator. Using the expression for the Hamiltonian

$$H = \frac{P^2}{2m} + \frac{1}{2}m\omega^2 X^2, \tag{46}$$

and the representation for X and P, we obtain the following form for the stationary state Schrödinger equation:

$$\frac{d^2\psi(p)}{dp^2} + \frac{2\beta p}{1-\beta p^2}\frac{d\psi(p)}{dp} + \left(1-\beta p^2\right)^2 \left(\epsilon - \eta^2 p^2\right)\psi(p) = 0, \tag{47}$$

where $-1/\sqrt{\beta} \leq p \leq 1/\sqrt{\beta}$ and

$$\epsilon = \frac{2E}{m\hbar^2\omega^2}, \qquad \eta = \frac{1}{m\hbar\omega}. \tag{48}$$

A. The quantum mechanical solution

Using the dimensionless variable $u = \sqrt{\beta}p$, Eq. (47) can be written as

$$\frac{d^2\psi(u)}{du^2} + \frac{2u}{1-u^2}\frac{d\psi(u)}{du} + \left(1-u^2\right)^2 \left(\epsilon' - \eta'^2 u^2\right)\psi(u) = 0, \tag{49}$$

where $-1 \leq u \leq 1$ and

$$\epsilon' = \frac{\epsilon}{\beta}, \qquad \eta' = \frac{\eta}{\beta}. \tag{50}$$

Now by changing the variable to $x = u - (1/3)u^3$ we have

$$-\frac{d^2\psi(x)}{dx^2} + \eta'^2\, V(x)\psi(x) = \epsilon'\psi(x), \tag{51}$$

where $-2/3 \leq x \leq 2/3$ and

$$V(x) = \left[\frac{1 - i\sqrt{3} + (-2)^{1/3}\left(3x + \sqrt{9x^2-4}\right)^{2/3}}{2^{2/3}\left(3x+\sqrt{9x^2-4}\right)^{1/3}}\right]^2, \tag{52}$$

is the effective potential (see Fig. 3). The boundary condition now is

$$\psi(x)\Big|_{\pm\frac{2}{3}} = 0. \tag{53}$$

To solve Eq. (51), we can expand the wave function in terms of the particle in a box eigenfunctions. Since the potential term $V(x)$ is an even function of x, to avoid large matrices, we use

$$\phi_m^{\mathrm{e}}(x) = \sqrt{\frac{1}{L}}\cos\left[\left(m-\frac{1}{2}\right)\frac{\pi x}{L}\right], \tag{54}$$

and

$$\phi_m^{\mathrm{o}}(x) = \sqrt{\frac{1}{L}}\sin\left(\frac{m\pi x}{L}\right), \tag{55}$$



FIG. 3. The effective potential $V(x)$ (blue line) and the ground state wave function (red line) for $\eta' = 100$.

basis functions ($m = 1, 2, \ldots$) for even and odd parity solutions, respectively, and write the wave function as $\psi(x) = \sum_m A_m \phi_m(x)$ which vanishes at $\pm L$. Now the boundary condition (53) reads $L = 2/3$.

The approximate solutions are the eigenvalues and the eigenfunctions of the ($N \times N$) Hamiltonian matrix $\mathbf{H}_N$ in the form

$$H_{mn} = \left(m - \frac{1}{2}\right)^2 \frac{\pi^2}{L^2} \delta_{mn} + D_{mn}^{\text{even}},$$

and

$$H_{mn} = \frac{m^2 \pi^2}{L^2} \delta_{mn} + D_{mn}^{\text{odd}}, \tag{56}$$

for even and odd states, respectively. Here, δ_{mn} is the kronecker's delta and

$$D_{mn}^{\text{even}} = \frac{\eta'^2}{L} \int_{-L}^{L} V(x) \cos\left[\left(m - \frac{1}{2}\right)\frac{\pi x}{L}\right] \cos\left[\left(n - \frac{1}{2}\right)\frac{\pi x}{L}\right] \mathrm{d}x, \tag{57}$$

$$D_{mn}^{\text{odd}} = \frac{\eta'^2}{L} \int_{-L}^{L} V(x) \sin\left(\frac{m\pi x}{L}\right) \sin\left(\frac{n\pi x}{L}\right) \mathrm{d}x, \tag{58}$$

where m and n run from 1 to N. In the usual diagonalization scheme with the particle in box basis functions, we need to adjust the domain L with respect to the number of basis functions in such way that the total error to be minimized [36]. However, for our case, since the boundary condition (53) has fixed the domain, i.e., $L = 2/3$, the accuracy of the solutions grows as the number of the basis increases. In Table II we have reported the first ten energy eigenvalues of the harmonic oscillator in the GUP* framework. Indeed $N = 30$ basis functions suffices to obtain nearly accurate results for the low lying energy eigenstates.

TABLE II. The energy eigenvalues of the harmonic oscillator in the GUP* framework. Here $\mathcal{E}_n = \epsilon'_n/\eta' = \epsilon_n/\eta = 2E_n/\hbar\omega$, $N = 30$, and $\eta' = 100$.

n	$\mathcal{E}_n^{\beta=0}$	$\mathcal{E}_n^{(SC)}$	$\mathcal{E}_n$	$\frac{\lvert\mathcal{E}_n-\mathcal{E}_n^{(SC)}\rvert}{\mathcal{E}_n}$
0	1	1.00251	1.00509	2.6×10^{-3}
1	3	3.02284	3.02559	9.1×10^{-4}
2	5	5.06411	5.06704	5.8×10^{-4}
3	7	7.12698	7.13011	4.4×10^{-4}
4	9	9.21216	9.21550	3.6×10^{-4}
5	11	11.3204	11.3240	3.2×10^{-4}
6	13	13.4524	13.4563	2.9×10^{-4}
7	15	15.6091	15.6133	2.7×10^{-4}
8	17	17.7913	17.7958	2.5×10^{-4}
9	19	20.0000	20.0049	2.4×10^{-4}

B. The semiclassical solution

The total energy in terms of ordinary variables is

$$E = \frac{p^2}{2m} + \frac{m\omega^2 x^2}{2\left(1-\beta p^2\right)^2}. \tag{59}$$

To find the approximate energy eigenvalues of the above Hamiltonian, we use the Wilson-Sommerfeld quantization rule in the form

$$\oint x\,\mathrm{d}p = \left(n+\frac{1}{2}\right)h, \qquad n = 0, 1, 2, \ldots, \tag{60}$$

where we have used $\oint \mathrm{d}(xp) = 0 = \oint x\,\mathrm{d}p + \oint p\,\mathrm{d}x$. This integral can be written as

$$\oint x\,\mathrm{d}p = \frac{2}{m\omega}\int_{-z}^{z}\left(1-\beta p^2\right)\sqrt{z^2-p^2}\,\mathrm{d}p, \tag{61}$$

where $z = \sqrt{2mE}$. So the semiclassical energy spectrum is given by

$$\begin{aligned} E_n^{(SC)} &= \frac{1-\sqrt{1-2m\beta\hbar\omega\left(n+\frac{1}{2}\right)}}{m\beta}, & (62)\\ &= -\frac{1}{8}\gamma\hbar\omega + \hbar\omega\left(n+\frac{1}{2}\right)\left(1+\frac{\gamma}{2}\right) + \frac{1}{2}\gamma\hbar\omega n^2 + \frac{1}{2}\gamma^2\hbar\omega\left(n+\frac{1}{2}\right)^3 + \mathcal{O}(\gamma^3), & (63)\end{aligned}$$



FIG. 4. Comparing semiclassical harmonic oscillator energy spectrum for KMM GUP (red line) and GUP* (blue line). We set $\gamma = 0.01$.

where $\gamma = \beta m \hbar \omega = \eta'^{-1}$. As it is shown in the appendix, the first three terms are similar to the energy spectrum of the harmonic oscillator in the KMM framework. In Fig. 4, we have depicted the energy spectrum in both $\mathrm{GUP}^{\mathrm{KMM}}$ and GUP* frameworks. Note that, in GUP* scenario, the energy is also bounded from above. Indeed, the maximum possible energy for the harmonic oscillator is

$$E_{max}^{(SC)} = \frac{1}{m\beta}, \tag{64}$$

and the number of states ($N = n + 1$) is finite, namely

$$n_{max} = \left\lfloor \frac{1}{2\gamma} - \frac{1}{2} \right\rfloor, \tag{65}$$

where $\lfloor x \rfloor$ denotes the largest integer not greater than x. So, to have at least one state, we should have

$$\gamma \leq 1. \tag{66}$$

As Fig. 4 shows, we have 50 states for $\gamma = 0.01$. The first ten semiclassical energy eigenvalues are presented in Table II [37]. As the table shows the semiclassical results agree well with the quantum mechanical energy spectrum. In fact, the relative error is less than 3×10^{-3} even for the ground state.

C. The classical solution

In the classical domain, the equations of motion are

$$\dot{X} = \{X, H\} = \frac{P}{m\,(1-\beta P^2)}, \tag{67}$$

$$\dot{P} = \{P, H\} = -\frac{m\omega^2 X}{1-\beta P^2}. \tag{68}$$

The solutions to these equations are

$$\omega t = \left(1-\frac{\epsilon}{2}\right)\arccos\left(\frac{P(t)}{P_{max}}\right) - \frac{\beta}{2}P(t)\sqrt{P_{max}^2 - P^2(t)}, \tag{69}$$

$$X(t) = -\frac{1-\beta P^2(t)}{m\omega^2}\frac{\mathrm{d}P(t)}{\mathrm{d}t}, \tag{70}$$

where

$$\epsilon = 2m\beta E, \qquad P_{max} = \sqrt{2mE}. \tag{71}$$

To first-order in β we have

$$P(t) = P_{max}\left(\cos\left[\left(1+\frac{\epsilon}{2}\right)\omega t\right] - \frac{\epsilon}{2}\sin^2\omega t\cos\omega t\right), \tag{72}$$

$$X(t) = X_{max}\left[\left(1+\frac{\epsilon}{2}\right)\cos\left[\left(1+\frac{\epsilon}{2}\right)\omega t\right] - \frac{\epsilon}{2}\sin^3\omega t\right], \tag{73}$$

where $X_{max} = \sqrt{2E/m\omega^2}$. As we have expected these results agree with the KMM proposal to $\mathcal{O}(\beta)$ [38].

It is straightforward to show that the infinitesimal phase space volume between equal energy contours E and $E + \mathrm{d}E$, and equal time contours t and $t + \mathrm{d}t$ can be written as

$$\mathrm{d}E\,\mathrm{d}t = (1-\beta P^2)\mathrm{d}X\,\mathrm{d}P. \tag{74}$$

Now, since by definition the left hand side of this equation is time independent, the right hand is also time independent.

V. A FORMALLY SELF-ADJOINT REPRESENTATION

Although the set of Eqs. (17) and (18) is an exact representation of the algebra presented in Eq. (16), it is not self-adjoint and does not preserve the ordinary nature of the position operator. To solve these difficulties, we can write $P = f(p)$ and preserve the ordinary nature of the position operator, i.e., $X = x$. Thus, using Eq. (16) we find $\frac{df}{dp} = \frac{1}{1-\beta f^2}$ which results in

$$f(p) - \frac{1}{3}\beta f^3(p) = p. \tag{75}$$

Consequently, the alternative representation in exact and perturbative form can be written as

$$\begin{aligned} X &= x, && (76) \\ P &= \frac{1 - i\sqrt{3} + (-2\beta)^{1/3}\left(3p + \sqrt{9p^2 - 4/\beta}\right)^{2/3}}{(2\beta)^{2/3}\left(3p + \sqrt{9p^2 - 4/\beta}\right)^{1/3}}, && (77) \\ &= p + \frac{1}{3}\beta p^3 + \frac{1}{3}\beta^2 p^5 + \frac{4}{9}\beta^3 p^7 + \cdots . && (78) \end{aligned}$$

Note that this representation is formally self-adjoint, i.e., $A = A^\dagger$ for $A \in \{X, P\}$. Also Eq. (77) shows that the momentum P has a upper bound

$$P_{max} = \frac{1}{\sqrt{\beta}}, \tag{79}$$

at $p = \frac{2}{3\sqrt{\beta}}$. Now X and P are symmetric operator on the dense domain S_∞ with respect to the following scalar product in the momentum space:

$$\langle\psi|\phi\rangle = \int_{-\frac{2}{3\sqrt{\beta}}}^{+\frac{2}{3\sqrt{\beta}}} \psi^*(p)\phi(p)\,\mathrm{d}p. \tag{80}$$

We have schematically depicted the behavior of P versus p in Fig 5.

In this representation, to write the Hamiltonian, it is more appropriate to use Eq. (78) and express the Hamiltonian perturbatively as

$$H = \frac{p^2}{2m} + V(x) + \beta\frac{p^4}{3m} + \beta^2\frac{7p^6}{18m} + \beta^3\frac{5p^8}{9m} + \mathcal{O}(\beta^4), \tag{81}$$

which agrees with perturbative version of the KMM proposal to $\mathcal{O}(\beta)$ [23]. In the quantum domain, this Hamiltonian results in the following generalized Schrödinger equation in quasiposition representation:

$$-\frac{\hbar^2}{2m}\frac{\partial^2\psi(x)}{\partial x^2} + \frac{\beta}{3m}\frac{\partial^4\psi(x)}{\partial x^4} - \frac{7\beta^2}{18m}\frac{\partial^6\psi(x)}{\partial x^6} + \frac{5\beta^3}{9m}\frac{\partial^8\psi(x)}{\partial x^8} + \mathcal{O}(\beta^4) + V(x)\psi(x) = E\,\psi(x), \tag{82}$$

where the extra terms are due to the GUP-corrected terms in Eq. (81). Notice Eq. (77) shows that this representation is naturally perturbative that is apparent from Eq. (82).



FIG. 5. Schematic behavior of P versus p in the second representation for the ordinary quantum mechanics (red line) and GUP* (blue line).

Note that for an operator A which is "formally" self-adjoint ($A = A^\dagger$) such as (76) and (78), this does not prove that A is truly self-adjoint because in general the domains $\mathcal{D}(A)$ and $\mathcal{D}(A^\dagger)$ may be different. The operator A with dense domain $\mathcal{D}(A)$ is said to be self-adjoint if $\mathcal{D}(A) = \mathcal{D}(A^\dagger)$ and $A = A^\dagger$. For instance, in this representation, X (76) is merely symmetric but not self-adjoint. To see this note that in this representation and in the momentum space the wave function $\phi(p)$ have to vanish at the end of the p interval $(-2/3\sqrt{\beta} < p < 2/3\sqrt{\beta})$, because the tangent function diverges there. So X is a derivative operator $i\hbar\partial/\partial p$ on an interval with Dirichlet boundary conditions. But this means that X cannot be self-adjoint because all candidates for the eigenfunctions of X, (the plane waves, which are even normalizable) are not in the domain of X because they do not obey Dirichlet boundary conditions. Calculating the domain of the adjoint of X shows that it is larger than that of X, so X is indeed not self-adjoint, i.e.,

$$\int_{-\frac{2}{3\sqrt{\beta}}}^{+\frac{2}{3\sqrt{\beta}}} \mathrm{d}p\, \psi^*(p) \left(i\hbar\frac{\partial}{\partial p}\right) \phi(p) = \int_{-\frac{2}{3\sqrt{\beta}}}^{+\frac{2}{3\sqrt{\beta}}} \mathrm{d}p\, \left(i\hbar\frac{\partial\psi(p)}{\partial p}\right)^* \phi(p) + i\hbar\,\psi^*(p)\phi(p)\Big|_{p=+\frac{2}{3\sqrt{\beta}}} - i\hbar\,\psi^*(p)\phi(p)\Big|_{p=-\frac{2}{3\sqrt{\beta}}}. \tag{83}$$

Now since $\phi(p)$ vanishes at $p = \pm\dfrac{2}{3\sqrt{\beta}}$, $\psi^*(p)$ can take any arbitrary value at the boundaries. The above equation implies that X is symmetric, but it is not a self-adjoint operator.

Although its adjoint $X^\dagger = i\hbar\partial/\partial p$ has the same formal expression, it acts on a different space of functions, namely

$$\mathcal{D}(X) = \left\{\phi, \phi' \in \mathcal{L}^2\left(\frac{-2}{3\sqrt{\beta}}, \frac{+2}{3\sqrt{\beta}}\right);\phi\left(\frac{+2}{3\sqrt{\beta}}\right) = \phi\left(\frac{-2}{3\sqrt{\beta}}\right) = 0\right\}, \tag{84}$$

$$\mathcal{D}(X^\dagger) = \left\{\psi, \psi' \in \mathcal{L}^2\left(\frac{-2}{3\sqrt{\beta}}, \frac{+2}{3\sqrt{\beta}}\right);\text{no other restriction on } \psi\right\}. \tag{85}$$

On the other hand, since there is no Dirichlet boundary conditions on the wave functions in the position space ($-\infty < x < \infty$), P is still self-adjoint.

To check the self-adjointness property of X, we can also use the the von Neumann's theorem [39, 40]. First we need to find the wave functions that satisfy the eigenvalue equation

$$X^\dagger\phi_\pm(p) = i\hbar\partial_p\phi_\pm(p) = \pm i\lambda\phi_\pm(x), \tag{86}$$

that reads

$$\phi_\pm(p) = \mathcal{C}_\pm e^{\mp\lambda p}. \tag{87}$$

Since both $\phi_\pm(p)$ belong to $\mathcal{L}^2\left(\frac{-2}{3\sqrt{\beta}}, \frac{+2}{3\sqrt{\beta}}\right)$, the deficiency indices are $(1, 1)$. Therefore, the position operator is not self-adjoint but has a one-parameter family of self-adjoint extensions which is in agreement with the previous result.

A. Free particle

In ordinary quantum mechanics, the free particle wave function $u_p(x)$ is defined as the eigenfunction of the momentum operator, namely

$$\hat{P}u_p(x) = p\, u_p(x), \tag{88}$$

where p is the eigenvalue. The momentum operator has the following representation in the position space

$$\hat{P} = \frac{\hbar}{i}\frac{\partial}{\partial x}. \tag{89}$$

So, from Eq. (88) we have

$$\frac{\hbar}{i}\frac{\partial u_p(x)}{\partial x} = p\, u_p(x), \tag{90}$$

which has the following solution

$$u_p(x) = \frac{1}{\sqrt{2\pi\hbar}} \exp\left(\frac{ipx}{\hbar}\right), \tag{91}$$

where the constant of integration is chosen to satisfy

$$\int_{-\infty}^{\infty} u_p^*(x) u_p(x') \mathrm{d}p = \delta(x - x'). \tag{92}$$

In the GUP scenario, to find the momentum eigenfunction in the quasiposition space, we write the eigenvalue equation as

$$\frac{1 - i\sqrt{3} + (-2\beta)^{1/3}\left(-3i\hbar\partial_\xi + \sqrt{-9\hbar^2\partial_\xi^2 - 4/\beta}\right)^{2/3}}{(2\beta)^{2/3}\left(-3i\hbar\partial_\xi + \sqrt{-9\hbar^2\partial_\xi^2 - 4/\beta}\right)^{1/3}} u_\wp(\xi) = \wp\, u_\wp(\xi), \tag{93}$$

where $\wp$ is the eigenvalue of P. Now, let us take the solution in the form of Eq. (91) and (93) at the same time, i.e.,

$$u_\wp(\xi) = \frac{1}{\sqrt{2\pi\hbar}} \exp\left(\frac{ip\,\xi}{\hbar}\right), \tag{94}$$

where $p = f(\wp)$. Inserting this solution in Eq. (93) results in

$$\frac{1 - i\sqrt{3} + (-2\beta)^{1/3}\left(3p + \sqrt{9p^2 - 4/\beta}\right)^{2/3}}{(2\beta)^{2/3}\left(3p + \sqrt{9p^2 - 4/\beta}\right)^{1/3}} = \wp, \tag{95}$$

or

$$p = \wp - \frac{\beta}{3}\wp^3, \tag{96}$$

so we have

$$u_\wp(\xi) = \frac{1}{\sqrt{2\pi\hbar}} \exp\left[\frac{i}{\hbar}\left(\wp - \frac{\beta}{3}\wp^3\right)\xi\right]. \tag{97}$$

The momentum eigenfunctions now satisfy

$$\int_{-\frac{2}{3\sqrt{\beta}}}^{+\frac{2}{3\sqrt{\beta}}} u_\wp^*(\xi') u_\wp(\xi) \mathrm{d}p = \int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \left(1 - \beta\wp^2\right) u_\wp^*(\xi') u_\wp(\xi) \mathrm{d}\wp, \tag{98}$$

$$= \frac{\hbar\sqrt{\beta}}{\pi(\xi - \xi')} \sin\left(\frac{2(\xi - \xi')}{3\hbar\sqrt{\beta}}\right), \tag{99}$$

in agreement with Eq. (28). Finally, the plain wave (97) has the maximum energy $E_{max} = 3/(2m\beta)$ which agrees with Eq. (43).

B. Particle in a box

As another application, let us consider a particle with mass m confined in an infinite one-dimensional box with length L

$$V(x) = \begin{cases} 0 & 0 < x < L, \\ \infty & \text{elsewhere.} \end{cases} \tag{100}$$

The corresponding eigenfunctions should satisfy the following generalized Schrödinger equation

$$-\frac{\hbar^2}{2m}\frac{\partial^2\psi_n(x)}{\partial x^2} + \frac{\beta\hbar^4}{3m}\frac{\partial^4\psi(x)}{\partial x^4} - \frac{7\beta^2\hbar^6}{18m}\frac{\partial^6\psi(x)}{\partial x^6} + \frac{5\beta^3\hbar^8}{9m}\frac{\partial^8\psi(x)}{\partial x^8} + \mathcal{O}(\beta^4) = E_n\,\psi_n(x), \tag{101}$$

for $0 < x < L$ and they also meet the boundary conditions $\psi_n(0) = \psi_n(L) = 0$. In Refs. [16, 22], the above equation is thoroughly solved to $\mathcal{O}(\beta)$ and its exact eigenvalues and eigenfunctions are found. Because of the boundary conditions, the eigenfunctions do not change with respect to the absence of the GUP ($\beta = 0$). However, the solutions exhibit the effect of GUP on the eigenvalues which are linear in GUP parameter [22]. These facts lead us to consider the following additional condition for eigenfunctions

$$-\frac{\hbar^2}{2m}\frac{\partial^2\psi_n(x)}{\partial x^2} = \varepsilon_n\psi_n(x), \qquad 0 < x < L, \tag{102}$$

where $\varepsilon_n = \frac{n^2\pi^2\hbar^2}{2mL^2}$. If this condition is also satisfied, we can write the second term in Eq. (101) in terms of $\psi_n(x)$, i.e.,

$$\frac{\partial^{2j}\psi_n(x)}{\partial x^{2j}} = \frac{-2m\varepsilon_n}{\hbar^2}\frac{\partial^{2(j-1)}\psi_n(x)}{\partial x^{2(j-1)}} = \cdots = \left(\frac{-2m\varepsilon_n}{\hbar^2}\right)^j \psi_n(x). \tag{103}$$

So, we have

$$H\psi_n(x) = \left(\varepsilon_n + \frac{4}{3}\beta m\varepsilon_n^2 + \frac{28}{9}\beta^2 m^2\varepsilon_n^3 + \frac{80}{9}\beta^3 m^3\varepsilon_n^4 + \cdots\right)\psi_n(x). \tag{104}$$

Now, comparing Eqs. (101) and (104) shows that

$$E_n = \varepsilon_n + \frac{4}{3}\beta m\varepsilon_n^2 + \frac{28}{9}\beta^2 m^2\varepsilon_n^3 + \frac{80}{9}\beta^3 m^3\varepsilon_n^4 + \cdots, \tag{105}$$

$$= \varepsilon_n \left[\frac{1 - i\sqrt{3} + (-2)^{1/3}\left(3\gamma_n + \sqrt{9\gamma_n^2 - 4}\right)^{2/3}}{4^{1/3}\left(3\gamma_n + \sqrt{9\gamma_n^2 - 4}\right)^{1/3}}\right]^2, \tag{106}$$

where $\gamma_n = 2\beta m\varepsilon_n$. To first order of GUP parameter we have $E_n = \frac{n^2\pi^2\hbar^2}{2mL^2} + \frac{\beta n^4\pi^4\hbar^4}{3mL^4}$ which is in agreement with the result of Ref. [22]. Moreover, because of Eq. (102) and the boundary conditions, the normalized eigenfunctions are $\psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right)$ [22]. These results show that in GUP* scenario there is no change in the particle in a box eigenfunctions but there is a positive shift in the energy spectrum which is proportional to the powers of β.

We now estimate the energy spectrum using the semiclassical scheme. For the particle in a box, the Wilson-Sommerfeld formula

$$\oint p \, \mathrm{d}x = nh, \qquad n = 1, 2, \cdots, \tag{107}$$

results in

$$p_n = \frac{nh}{L}. \tag{108}$$

Since the high energy momentum P depends on the low energy momentum through $p_n = P_n - (1/3)\beta P_n^3$ (75), the semiclassical energy spectrum is given by

$$\begin{aligned} E_n^{(SC)} &= \frac{P_n^2}{2m}, \\ &= \left[\frac{1 - i\sqrt{3} + (-2\beta)^{1/3}\left(3p_n + \sqrt{9p_n^2 - 4/\beta}\right)^{2/3}}{\sqrt{2m}(2\beta)^{2/3}\left(3p_n + \sqrt{9p_n^2 - 4/\beta}\right)^{1/3}}\right]^2. \end{aligned} \tag{109}$$

It is straightforward to check that the semiclassical result (109) exactly coincide with the quantum mechanical spectrum (105) as well as in ordinary quantum mechanics. Therefore, the maximal energy of a particle in a box reads

$$E_{max} = \frac{1}{2m\beta}. \tag{110}$$

C. WKB approximation

To check the validity of the Wilson-Sommerfeld quantization rule for this modified quantum mechanics, we need to show that the zeroth-order wave function, which satisfies the generalized Schrödinger equation (82), can be written as $\psi(x) \simeq \exp\left[(i/\hbar)\int p \, \mathrm{d}x\right]$. So let us take

$$\psi(x) = e^{i\Phi(x)}, \tag{111}$$

where $\Phi(x)$ can be expanded as a power series in $\hbar$ in the semiclassical approximation, i.e.,

$$\Phi(x) = \frac{1}{\hbar}\sum_{n=0}^{\infty} \hbar^n \Phi_n(x). \tag{112}$$

So we have

$$\frac{\partial^2\psi(x)}{\partial x^2} = \left(-\Phi'^2 + i\Phi''\right)\psi(x), \tag{113}$$

$$\frac{\partial^4\psi(x)}{\partial x^4} = \left(\Phi'^4 - 6i\Phi'^2\Phi'' - 3\Phi''^2 - 4\Phi'''\Phi' + i\Phi''''\right)\psi(x), \tag{114}$$

$$\vdots \qquad\qquad \vdots$$

where Φ' indicates the derivative of Φ with respect to x. Now to zeroth-order ($\Phi(x) \simeq \Phi_0(x)/\hbar$) and for $\hbar \to 0$ we obtain

$$\Phi_0'^2 + \frac{2}{3}\beta\Phi_0'^4 + \frac{7}{9}\beta^2\Phi_0'^6 + \mathcal{O}(\beta^3) = 2m\left(E - V(x)\right). \tag{115}$$

Now the comparison with Eq. (81) shows $\Phi_0' = p$ and consequently

$$\psi(x) \simeq \exp\left[\frac{i}{\hbar}\int p\,\mathrm{d}x\right], \tag{116}$$

which is the usual zeroth-order WKB wave function obeying the Wilson-Sommerfeld quantization rule.

VI. GENERALIZATION TO D DIMENSIONS

We now extend the developed formalism in Sec. II to D spatial dimensions. We then present the generalized Poisson brackets in the classical limit and study the density of states.

A. Generalized Heisenberg algebra for D dimensions

A natural generalization of the one-dimensional commutation relation (16) that preserves the rotational symmetry is

$$[X_i, P_j] = \frac{i\hbar\delta_{ij}}{1-\beta P^2}, \tag{117}$$

where $P^2 = \sum_{i=1}^{D} P_i P_i$. This relation implies a nonzero minimal uncertainty and a maximal observable momentum in each position coordinate. If the components of the momentum operator are assumed to be commutative

$$[P_i, P_j] = 0, \tag{118}$$

then the Jacobi identity determines the commutation relations between the components of the position operator as

$$[X_i, X_j] = \frac{2i\hbar\beta}{\left(1-\beta P^2\right)^2}\left(P_i X_j - P_j X_i\right), \tag{119}$$

which results in a noncommutative geometric generalization of position space. To exactly satisfy these commutation relations, the position and momentum operators in the momentum space representation can be written as

$$P_i\phi(p) = p_i\phi(p), \tag{120}$$

$$X_i\phi(p) = \frac{i\hbar}{1-\beta p^2}\partial_{p_i}\phi(p). \tag{121}$$

X_i and P_j are now symmetric operator on the domain S_∞ with respect to the scalar product:

$$\langle\psi|\phi\rangle = \int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \mathrm{d}^{\mathrm{D}}p\left(1-\beta p^2\right)\psi^*(p)\phi(p), \tag{122}$$

where $p^2 = \sum_{i=1}^{D} p_i p_i$. The identity operator is

$$1 = \int_{-1/\sqrt{\beta}}^{+1/\sqrt{\beta}} \frac{\mathrm{d}^{\mathrm{D}}p}{\left(1-\beta p^2\right)}|p\rangle\langle p|, \tag{123}$$

and the scalar product of momentum eigenstates is

$$\langle p|p'\rangle = \frac{\delta^D(p-p')}{1-\beta p^2}. \tag{124}$$

In this representation, the components of the momentum operator are still essentially self-adjoint, however the components of the position operators are merely symmetric and do not have physical eigenstates.

Since the commutation relations (117)–(119) do not break the rotational symmetry, we can express the generators of rotations in terms of the position and momentum operators as

$$L_{ij} \equiv \left(1-\beta P^2\right)\left(X_i P_j - X_j P_i\right), \tag{125}$$

as the generalization of the ordinary orbital angular momentum. Now the momentum space representation of the generators of rotations is

$$L_{ij}\psi(p) = -i\hbar\left(p_i\partial_{p_j} - p_j\partial_{p_i}\right)\psi(p), \tag{126}$$

and

$$[P_i, L_{jk}] = i\hbar\left(\delta_{ik}P_j - \delta_{ij}P_k\right), \tag{127}$$

$$[X_i, L_{jk}] = i\hbar \left(\delta_{ik} X_j - \delta_{ij} X_k\right), \tag{128}$$

$$[L_{ij}, L_{kl}] = i\hbar \left(\delta_{ik} L_{jl} - \delta_{il} L_{jk} + \delta_{jl} L_{ik} - \delta_{jk} L_{il}\right), \tag{129}$$

as well as in ordinary quantum mechanics. However, the geometry is noncommutative, namely

$$[X_i, X_j] = \frac{-2i\hbar\beta}{\left(1-\beta P^2\right)^2} L_{ij}. \tag{130}$$

B. Density of states

The right hand side of Eq. (16) shows that the "effective" value of $\hbar$ is P dependent. So the size of the unit cell in the phase space that is occupied by each quantum state can be also considered of as being momentum dependent. This fact changes the momentum dependence of the density of states and affect the calculation of the cosmological constant, the blackbody radiation spectrum, etc. Similar to the KMM algebra [38], we should first check that any volume of the phase space evolves such that the number of states inside it does not change with respect to time as the analog of the Liouville theorem.

The Poisson brackets in classical mechanics correspond quantum mechanical commutators via

$$\frac{1}{i\hbar}[A, B] \Longrightarrow \{A, B\}. \tag{131}$$

Thus the classical limits of Eqs. (117)–(119) are given by

$$\{X_i, P_j\} = \frac{\delta_{ij}}{\left(1-\beta P^2\right)}, \tag{132}$$

$$\{P_i, P_j\} = 0, \tag{133}$$

$$\{X_i, X_j\} = \frac{2\beta}{\left(1-\beta P^2\right)^2}\left(P_i X_j - P_j X_i\right), \tag{134}$$

and the Heisenberg equations for the coordinates and momenta read (i, j run over the spatial dimensions and the summation convention is assumed)

$$\dot{X}_i = \{X_i, H\} = \{X_i, P_j\}\frac{\partial H}{\partial P_j} + \{X_i, X_j\}\frac{\partial H}{\partial X_j}, \tag{135}$$

$$\dot{P}_i = \{P_i, H\} = -\{X_j, P_i\}\frac{\partial H}{\partial X_j}. \tag{136}$$

We now prove that the weighted phase space volume

$$\left(1-\beta P^2\right)^D \mathrm{d}^D X\, \mathrm{d}^D P, \tag{137}$$

is invariant under time evolution as the analog of the Liouville theorem. The evolution of X_i and P_i during an infinitesimal time interval δt is

$$X_i' = X_i + \delta X_i, \tag{138}$$

$$P_i' = P_i + \delta P_i, \tag{139}$$

where

$$\delta X_i = \left[\{X_i, P_j\} \frac{\partial H}{\partial P_j} + \{X_i, X_j\} \frac{\partial H}{\partial X_j} \right] \delta t, \tag{140}$$

$$\delta P_i = \left[-\{X_j, P_i\} \frac{\partial H}{\partial X_j} \right] \delta t. \tag{141}$$

After this infinitesimal evolution, the infinitesimal phase space volume is changed according to

$$\mathrm{d}^D X' \, \mathrm{d}^D P' = \left| \frac{\partial(X_1', \cdots, X_D', P_1', \cdots, P_D')}{\partial(X_1, \cdots, X_D, P_1, \cdots, P_D)} \right| \mathrm{d}^D X \, \mathrm{d}^D P. \tag{142}$$

where

$$\begin{aligned} \frac{\partial X_i'}{\partial X_j} &= \delta_{ij} + \frac{\partial \delta X_i}{\partial X_j}, & \frac{\partial X_i'}{\partial P_j} &= \frac{\partial \delta X_i}{\partial P_j}, \\ \frac{\partial P_i'}{\partial X_j} &= \frac{\partial \delta P_i}{\partial X_j}, & \frac{\partial P_i'}{\partial P_j} &= \delta_{ij} + \frac{\partial \delta P_i}{\partial P_j}. \end{aligned} \tag{143}$$

The Jacobian can be calculated to first-order in δt as

$$\left| \frac{\partial(X_1', \cdots, X_D', P_1', \cdots, P_D')}{\partial(X_1, \cdots, X_D, P_1, \cdots, P_D)} \right| = 1 + \left(\frac{\partial \delta X_i}{\partial X_i} + \frac{\partial \delta P_i}{\partial P_i} \right) + \cdots . \tag{144}$$

So we have

$$\begin{aligned} \left(\frac{\partial \delta X_i}{\partial X_i} + \frac{\partial \delta P_i}{\partial P_i} \right) \frac{1}{\delta t} &= \frac{\partial}{\partial X_i} \left[\{X_i, P_j\} \frac{\partial H}{\partial P_j} + \{X_i, X_j\} \frac{\partial H}{\partial X_j} \right] - \frac{\partial}{\partial P_i} \left[\{X_j, P_i\} \frac{\partial H}{\partial X_j} \right], \\ &= \left[\frac{\partial}{\partial X_i} \{X_i, P_j\} \right] \frac{\partial H}{\partial P_j} + \{X_i, P_j\} \frac{\partial^2 H}{\partial X_i \partial P_j} + \left[\frac{\partial}{\partial X_i} \{X_i, X_j\} \right] \frac{\partial H}{\partial X_j} \\ &\quad + \{X_i, X_j\} \frac{\partial^2 H}{\partial X_i \partial X_j} - \left[\frac{\partial}{\partial P_i} \{X_j, P_i\} \right] \frac{\partial H}{\partial X_j} - \{X_j, P_i\} \frac{\partial^2 H}{\partial P_j \partial X_i}, \\ &= \left[\frac{\partial}{\partial X_i} \{X_i, X_j\} \right] \frac{\partial H}{\partial X_j} - \left[\frac{\partial}{\partial P_i} \{X_j, P_i\} \right] \frac{\partial H}{\partial X_j}, \\ &= \left[-\frac{2\beta(D-1)}{(1-\beta P^2)^2} P_j \right] \frac{\partial H}{\partial X_j} - \left[\frac{2\beta}{(1-\beta P^2)^2} P_j \right] \frac{\partial H}{\partial X_j}, \\ &= -\frac{2\beta D}{(1-\beta P^2)^2} P_j \frac{\partial H}{\partial X_j}, \end{aligned} \tag{145}$$

which to first-order in δt results in

$$\mathrm{d}^D X' \,\mathrm{d}^D P' = \mathrm{d}^D X \,\mathrm{d}^D P \left[1 - \frac{2\beta D}{\left(1-\beta P^2\right)^2} P_j \frac{\partial H}{\partial X_j} \delta t\right]. \tag{146}$$

Moreover

$$\begin{aligned}
1 - \beta P'^2 &= 1 - \beta (P_i + \delta P_i)^2, \\
&= 1 - \beta \left(P^2 + 2P_i \delta P_i + \cdots\right), \\
&= 1 - \beta \left(P^2 - 2P_i \{X_i, P_j\} \frac{\partial H}{\partial X_j} \delta t + \cdots\right), \\
&= 1 - \beta \left(P^2 - \frac{2P_i}{1-\beta P^2} \frac{\partial H}{\partial X_i} \delta t + \cdots\right), \\
&= (1-\beta P^2) + \frac{2\beta P_i}{1-\beta P^2} \frac{\partial H}{\partial X_i} \delta t + \cdots, \\
&= (1-\beta P^2) \left[1 + \frac{2\beta P_i}{\left(1-\beta P^2\right)^2} \frac{\partial H}{\partial X_i} \delta t + \cdots\right].
\end{aligned} \tag{147}$$

Therefore, to first-order in δt

$$\left(1 - \beta P'^2\right)^D = \left(1-\beta P^2\right)^D \left[1 + \frac{2\beta D}{\left(1-\beta P^2\right)^2} P_i \frac{\partial H}{\partial X_i} \delta t\right], \tag{148}$$

Now using Eqs. (146) and (148), it is obvious that the weighted phase space volume Eq. (137) is an invariant, i.e.,

$$\left(1 - \beta P'^2\right)^D \mathrm{d}^D X' \,\mathrm{d}^D P' = \left(1-\beta P^2\right)^D \mathrm{d}^D X \,\mathrm{d}^D P. \tag{149}$$

C. The cosmological constant

The cosmological constant can be obtained by summing over the zero-point energies of the harmonic oscillator's momentum states. Using the canonical form of the zero-point energy of each oscillator with mass m

$$\frac{1}{2}\hbar\omega = \frac{1}{2}\sqrt{p^2+m^2}, \tag{150}$$

the sum over all momentum states per unit volume is

$$\begin{aligned}
\Lambda(m) &= \int \mathrm{d}^3 p \left(1-\beta p^2\right)^3 \left(\frac{1}{2}\sqrt{p^2+m^2}\right), \\
&= 2\pi \int_0^{1/\sqrt{\beta}} \mathrm{d}p \left(1-\beta p^2\right)^3 p^2 \sqrt{p^2+m^2}, \\
&= \frac{\pi}{20\,\beta^2} f(\beta m^2)
\end{aligned} \tag{151}$$

where

$$f(x) = \frac{1}{96}\Big[\sqrt{1+x}(96 + 192x + 476x^2 + 380x^3 + 105x^4)$$
$$-(480x^2 + 720x^3 + 450x^4 + 105x^5)\cosh^{-1}\left(\sqrt{x}\right)\Big], \tag{152}$$

and $f(0) = 0$. In the massless limit we find

$$\Lambda(0) = \frac{\pi}{20\beta^2} = \frac{1}{10}\left[\Lambda(0)\right]^{\mathrm{KMM}}, \tag{153}$$

that is ten times smaller than the massless cosmological constant predicted by the KMM proposal [38]. This finite result is due to the vanishing of the density of states at high momenta where $p = 1/\sqrt{\beta}$ plays the role of the UV cutoff. So in this scenario we do not need to put by hand an arbitrary scale as the UV cutoff and the cosmological constant is automatically rendered finite. Note that since $1/\sqrt{\beta}$ is proportional to the Planck mass M_{Pl}, $\Lambda(0)$ is too large in practice and consequently the cosmological constant problem still remains unsolved. However, our formulation gives the better estimation of Λ with respect to that obtained in the KMM framework.

D. The blackbody radiation spectrum

Because of the weight factor $(1 - \beta P^2)^3$ in 3-dimensions, the average energy of the electromagnetic field per unit volume at temperature T is given by

$$\begin{aligned}\overline{E} &= \frac{8\pi}{c^3}\int_0^\infty \mathrm{d}\nu \left(1 - \beta\left(\frac{h\nu}{c}\right)^2\right)^3 \left(\frac{h\nu^3}{e^{h\nu/k_BT} - 1}\right), \\ &= \int_0^\infty \mathrm{d}\nu\, u_\beta(\nu, T),\end{aligned} \tag{154}$$

where

$$u_\beta(\nu, T) = \left(1 - \left(\frac{\nu}{\nu_\beta}\right)^2\right)^3 u_0(\nu, T). \tag{155}$$

Here

$$u_0(\nu, T) = \frac{8\pi h\nu^3}{c^3}\frac{1}{e^{h\nu/k_BT} - 1}, \tag{156}$$

is the ordinary spectrum function and $\nu_\beta = c/h\sqrt{\beta}$. To show the effect of the minimal length uncertainty and the maximal momentum on the shape of the spectral function, we



FIG. 6. The blackbody radiation spectrum in the GUP* framework at temperature $T = 0.1\,T_\beta$.

have depicted the functions

$$f_0(\nu,T) = \frac{(\nu/\nu_\beta)^3}{e^{(\nu/\nu_\beta)(T_\beta/T)} - 1}, \tag{157}$$

$$f_\beta(\nu,T) = \left(1 - (\nu/\nu_\beta)^2\right)^3 \, f_0(\nu,T), \tag{158}$$

in Figs. 6 and 7, and compared them with the case of just the minimal length uncertainty [38]

$$f_\beta^{\mathrm{KMM}}(\nu,T) = \frac{1}{\left(1 + (\nu/\nu_\beta)^2\right)^3}\, f_0(\nu,T), \tag{159}$$

where $T_\beta = c/k_B\sqrt{\beta}$. As the figure shows, for small frequencies ($\nu \ll \nu_\beta$), $f_\beta(\nu,T)$ closely coincides with f_β^{KMM}. However, it deviates from f_β^{KMM} as the frequency increases.

VII. CONCLUSIONS

In this paper, we have presented a higher order generalized uncertainty principle that implies both a minimal length uncertainty and a maximal momentum proportional to $\hbar\sqrt{\beta}$ and $1/\sqrt{\beta}$, respectively. The first exact representation (7) and (8) was not self-adjoint, but preserves the ordinary nature of the momentum operator. We found the exact eigenfunctions of the position operator and quantum mechanical and semiclassical energy spectrum of the harmonic oscillator in this representation and showed that the energy spectrum is also bounded from above. The second representation (76) and (77) was formally self-adjoint and



FIG. 7. The blackbody radiation spectrum in the GUP* framework at temperature $T = T_\beta$.

resulted in the perturbative generalized Schrödinger equation in quasiposition space. We exactly studied the problems of the free particle and the particle in a box and showed that the existence of the maximal momentum $P_{max} = 1/\sqrt{\beta}$ is manifest from this representation. We then generalized our proposal to D dimensions and found the invariant density of states. We showed that the blackbody radiation spectrum will be modified at higher frequencies and compared our results with the KMM proposal. Although the cosmological constant was rendered finite, the smallness of the GUP parameter resulted in a large cosmological constant that could not solve the cosmological constant problem. However, our calculated cosmological constant was a better estimation with respect to the presence of just the minimal length. Here we implemented a momentum cutoff not through terms like P^2 on the right hand side of the commutation relations. Instead, we implemented the momentum cutoff through a function of P with a singularity. So the momentum space is cut into several sectors that decouple from each other. The sectors are separated from each other at the singularities of the function of P that is used. Technically, we have inequivalent irreducible representations of the commutation relations, one each in each sector [41]. This type of issue with the various sectors can be avoided, as it is indicated in Ref. [42]. Applied to our case, the trick would be to write the right hand side of the commutation relation not as a fraction but instead to expand it out as a geometric series. It has a finite radius of convergence and that rules out all representations beyond the singularity.

Appendix: Harmonic oscillator spectrum in the KMM framework

In the context of the KMM proposal, the total energy in terms of ordinary variables is given by

$$E = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 \left(1 + \beta p^2\right)^2 x^2. \tag{A.1}$$

Now the Wilson-Sommerfeld integral can be written as

$$\oint x \, \mathrm{d}p = \frac{2}{m\omega} \int_{-z}^{z} \frac{\sqrt{z^2 - p^2}}{1 + \beta p^2} \mathrm{d}p = \left(n + \frac{1}{2}\right) h, \tag{A.2}$$

where $z = \sqrt{2mE}$. So the semiclassical energy spectrum is given by

$$E_n^{(SC)} = -\frac{1}{8}\gamma\hbar\omega + \hbar\omega \left(n + \frac{1}{2}\right)\left(1 + \frac{\gamma}{2}\right) + \frac{1}{2}\gamma\hbar\omega n^2, \tag{A.3}$$

where $\gamma = \beta m\hbar\omega$. This result agrees (up to a constant) with the exact solution to first order of the GUP parameter [7]

$$\begin{aligned} E_n^{exact} &= \hbar\omega \left(n + \frac{1}{2}\right)\left(\sqrt{1 + \gamma^2/4} + \gamma/2\right) + \frac{1}{2}\gamma\hbar\omega n^2 \\ &= E_n^{(SC)} + \frac{1}{8}\gamma\hbar\omega + \mathcal{O}(\gamma^2), \end{aligned} \tag{A.4}$$

and gives the correct n^2 dependence behavior. An alternative derivation of Eq. (A.3) is also presented in Ref. [25].

ACKNOWLEDGMENTS

I am very grateful to Achim Kempf for fruitful discussions and suggestions.

---

[1] G. Veneziano, Europhys. Lett. **2**, 199 (1986).

[2] E. Witten, Phys. Today **49**, 24 (1996).

[3] D. Amati, M. Ciafaloni, G. Veneziano, Phys. Lett. B **216**, 41 (1989); Nucl. Phys. B **347**, 550 (1990); Nucl. Phys. B **403**, 707 (1993).

[4] K. Konishi, G. Paffuti, P. Provero, Phys. Lett. B **234**, 276 (1990).

[5] L.J. Garay, Int. J. Mod. Phys. A **10**, 145 (1995).

[6] M. Maggiore, Phys. Lett. B **319**, 83 (1993).

[7] A. Kempf, G. Mangano, R.B. Mann, Phys. Rev. D **52**, 1108 (1995).

[8] A. Kempf, G. Mangano, Phys. Rev. D **55**, 7909 (1997).

[9] M. Maggiore, Phys. Lett. B **304**, 65 (1993).

[10] F. Scardigli, Phys. Lett. B **452**, 39 (1999).

[11] M. Maggiore, Phys. Rev. D **49**, 5182 (1994), arXiv:hep-th/9305163.

[12] S. Hossenfelder et al., Phys. Lett. B **575**, 85 (2003), arXiv:hep-th/0305262.

[13] C. Bambi, F.R. Urban, Class. Quantum Grav. **25**, 095006 (2008), arXiv:0709.1965.

[14] K. Nozari and B. Fazlpour, Gen. Relativ. Grav. **38**, 1661 (2006).

[15] R. Banerjee and S. Ghosh, Phys. Lett. B **688**, 224 (2010), arXiv:1002.2302.

[16] P. Pedram, Int. J. Mod. Phys. D **19**, 2003 (2010).

[17] K. Nozari and P. Pedram, Europhys. Lett. **92**, 50013 (2010), arXiv:1011.5673.

[18] B. Vakili, Phys. Rev. D **77**, 044023 (2008).

[19] M.V. Battisti and G. Montani, Phys. Rev. D **77**, 023518 (2008).

[20] B. Vakili and H.R. Sepangi, Phys. Lett. B **651**, 79 (2007).

[21] M.V. Battisti and G. Montani, Phys. Lett. B **656**, 96 (2007).

[22] K. Nozari, T. Azizi, Gen. Relativ. Gravit. **38**, 735 (2006).

[23] S. Das and E.C. Vagenas, Phys. Rev. Lett. **101**, 221301 (2008), arXiv:0810.5333.

[24] K. Nouicer, Phys. Lett. B **646**, 63 (2007).

[25] P. Pedram, Phys. Rev. D **85**, 024016 (2012), arXiv:1112.2327.

[26] P. Pedram, Phys. Lett. B (2012), DOI:10.1016/j.physletb.2012.03.015.

[27] J. Magueijo and L. Smolin, Phys. Rev. Lett. **88**, 190403 (2002), arXiv:hep-th/0112090.

[28] J. Magueijo and L. Smolin, Phys. Rev. D **71**, 026010 (2005), arXiv:hep-th/0401087.

[29] J.L. Cortes and J. Gamboa, Phys. Rev. D **71**, 065015 (2005), arXiv:hep-th/0405285.

[30] P. Pedram, Europhys. Lett. **89**, 50008 (2010), arXiv:1003.2769.

[31] P. Pedram, K. Nozari, and S.H. Taheri, JHEP **03**, 093 (2011).

[32] A.F. Ali, S. Das, and E.C. Vagenas, Phys. Lett. B **678**, 497 (2009).

[33] S. Das, E.C. Vagenas, and A.F. Ali, Phys. Lett. B **690**, 407 (2010).

[34] A.F. Ali, S. Das, and E.C. Vagenas, Phys. Rev. D **84**, 044013 (2011).

[35] A. Kempf, arXiv:hep-th/9603115.

[36] P. Pedram, M. Mirzaei, and S.S. Gousheh, Mol. Phys. **108**, 1949 (2010).

[37] We have used the relation $\mathcal{E}_n^{(SC)} = 2\eta'\Big(1 - \sqrt{1 - 2\eta'^{-1}(n + 1/2)}\Big)$.

[38] L.N. Chang, D. Minic, N. Okamura, T. Takeuchi, Phys. Rev. D **65**, 125028 (2002).

[39] N.I. Akhiezer and I.M. Glazman, *Theory of Linear Operators in Hilbert Space* (Dover, New York, 1993).

[40] G. Bonneau, J. Faraut, and G. Valent, Am. J. Phys. **69**, 322 (2001).

[41] Private communication with Achim Kempf.

[42] A. Kempf, Phys. Rev. Lett. **92** (2004) 221301.